


FITC anti-mouse H-2Kd/H-2Dd

Catalog No.

FMFH2KdDd-01-025
 FMFH2KdDd-01-100

Size

25 Tests
 100 Tests



C57BL/6J splenocytes stained with FITC anti-mouse H-2Kd/H-2Dd (colored histogram) or blank control(gray histogram)

Clone:

4A34-1-2S

Isotype:

mouse IgG2a

Other Name

MHC class I

Storage Buffer:

Phosphate-buffered solution, pH 7.2-7.4, containing 0.1% Proclin 300 and 0.2%BSA

Applications:

FC

Vol.per.Test:

5µl/test

Excitation Laser

Blue Laser (488 nm)

Ex/Em Max

494nm/520nm

Reactivity:

Mouse

Storage Condition:

Store at 4° C. DO NOT FREEZE. LIGHT SENSITIVE MATERIAL.

Description:

The "classical" MHC Class I molecules are histocompatibility antigens encoded by the H-2 gene complex and consist of heterodimers of highly polymorphic α chains noncovalently associated with the invariant β 2-microglobulin. These antigens are expressed on most nucleated cells but expression varies on different cell types. MHC Class I molecules present endogenously synthesized peptides to CD8+ T lymphocytes, which are usually cytotoxic T cells. MHC Class I antigens expressed on thymic epithelial cells regulate the positive and negative selection of CD8+ T cells during T cell ontogeny. The monoclonal antibody 4A34-1-2S binds to a common determinant in the α 3 domains of H-2Kd and H-2Dd in the presence or absence of β 2 microglobulin. It cross reacts with the α 3 domain of H-2Kb.

Related Protocol:

4A Biotech Inc.

Add:7905 Silverton Avenue, Suite 117, San Diego, CA92126,USA

Website: <https://4abio.com/en/> E-mail: sales@4abio.com

Cell Surface Flow Cytometry Staining Protocol

Related Products:

Cat NO	Name
FXP001-100	RBC Lysing Buffer (10x)
FXP005	Cell Staining Buffer
FMU16/32-02-100	Purified anti-mouse CD16/32
FMCF002-01-100	FITC mouse IgG2a Isotype Control

REFERENCES:

1. Ozato K, Mayer NM, Sachs DH. Monoclonal antibodies to mouse major histocompatibility complex antigens. IV. A series of hybridoma clones producing anti-H-2d antibodies and an examination of expression of H-2d antigens on the surface of these cells. Transplantation. 1982;34:113-9. (Immunogen)
2. Burgert H, Maryanski JL, Kvist S. "E3/19K" protein of adenovirus type 2 inhibits lysis of cytolytic T lymphocytes by blocking cell-surface expression of histocompatibility class I antigens. Proc Natl Acad Sci USA. 1987;84:1356-60. (IP)
3. Lenz A, Heufler C, Rammensee H, Glassl H, Koch F, Romani N, et al. Murine epidermal Langerhans cells express significant amounts of class I major histocompatibility complex antigens. Proc Natl Acad Sci USA. 1989;86:7527-31. (FC, ICC, CMDC).

4A Biotech Inc.

Add:7905 Silverton Avenue, Suite 117, San Diego, CA92126,USA

Website: <https://4abio.com/en/> E-mail: sales@4abio.com